

Got Bunions? Get Lapiplasty® 3D Bunion Correction®

- Fixes the root cause (an unstable joint)^{1,2}
- Provides a 3D fix for a 3D problem
- Get back on your feet quickly* in a walking boot^{1,2}

Are Bunions Limiting Your Activities and Lifestyle?

Bunions can be very painful. With each step, your entire body weight rests on that bunion. Ouch! Conservative treatment options can help alleviate pain, but will not fix the problem because bunions are a progressive disorder that will not go away on their own without surgical correction.⁵

Bunions Are More Than a Bump

Bunion Surgery Has Changed

A common misconception is that a bunion is simply a "bump" that can be "shaved off." But, bunions are more than a bump – they are complex 3D problems caused by an unstable joint in the middle of the foot. While the majority of bunion surgery is only 2D and fails to address the root cause, the Lapiplasty® Procedure provides a 3D correction and secures the unstable joint.^{1, 2}



Fix It Right The First Time™


How Does Lapiplasty® 3D Bunion Correction® Work?

While traditional 2D osteotomy surgery cuts & shifts the bone to address the cosmetic bump, Lapiplasty® 3D Bunion Correction® does more — it corrects the entire bone in 3D and secures the unstable foundation to get you back on your feet typically within 3-10 days in a walking boot.^{3,4} The Lapiplasty® Procedure has also shown low recurrence; 97% and 99% maintain 3D correction in 13 and 17 months respectively.^{3,6}

[See How It Works](#)

Lapiplasty®


3D Bunion Correction



- ✓ Returns entire bone to normal alignment; a 3D correction³
- ✓ Secures the root cause; an unstable joint^{1,2}
- ✓ Get back on your feet quickly in a boot; many cases within 2 weeks^{3, 4}

Traditional

2D Bunion Surgery



- Unnaturally cuts & shifts bone; only a 2D correction
- Addresses cosmetic "bump" only; not the root cause
- 1 day to 6 weeks non-weight bearing in a post-operative shoe⁷

Get Back on Your Feet Quickly

Recovery From Lapiplasty®

A typical recovery timeline is detailed below. However, timepoints and doctor's orders vary based on each patient's condition. Be sure to ask your doctor what your particular recovery protocol will look like.



See the Results

Lapiplasty® Before & After Results

← Slide To See More →



Changing Bunion Correction for Good

The Positive Effect of Lapiplasty®



67 million

Americans are affected by a bunion deformity.⁸



of Lapiplasty® patients maintained their 3D bunion correction at 13 months.³



10.9

average days for Lapiplasty® patients to begin bearing weight on their operative foot in a boot.³




21

clinical publications directly support the Lapiplasty® procedure.

Real People, Real Stories

Hear What Patients Are Saying About Lapiplasty® 3D Bunion Correction®



Frequently Asked Questions

Is the Lapiplasty® Procedure covered by insurance?



Can the Lapiplasty® Procedure be performed as an outpatient procedure and how long does it last?



Can I wear my choice of footwear again?




When can I return to sports / activities?



Can the Lapiplasty® Procedure be performed on someone who has previously had traditional surgery?



Looking for More In-Depth Information Including Benefit and Risk?





Lapiplasty.com

A comprehensive patient resource for Lapiplasty® patients

Visit Lapiplasty.com →



Patient Risk Information

What are the risks associated with the Lapiplasty® Procedure?

Download Patient Risk Information →



Lapiplasty® Brochure

Learn more about Lapiplasty® in English or Spanish

Download in English or Spanish →

References

*typically within 3-10 days

1. Dayton P, et al. J Foot Ankle Surg. 2016. 55:567-71.
2. Dayton P, et al. J Foot Ankle Surg. 2018. 57:766-770.
3. Ray J, et al. Foot Ankle Int. 2019 Aug;40(8):955-960
4. Dayton P, et al. J Foot Ankle Surg. 2019. 58:427-433
5. American College of Foot and Ankle Surgeons (ACFAS) Website ©2023 <https://www.foothealthfacts.org/conditions/bunions>
6. Dayton P, et al. J Foot Ankle Surg. 2020, 59(2): 291-297
7. Ahn J, Lee HS, Seo JH, Kim JY. Foot & Ankle Int. 2016; 37(6):589-595
8. Nix S, et al. J Foot Ankle Res. 2010.

M2346A

Important Risk Information: Only a surgeon can tell if the Lapiplasty® Procedure is right for you. As with any medical treatment, individual results may vary. There are potential risks and recovery takes time. For more information about recovery from the Lapiplasty® Procedure, see the **recovery information** and discuss the post-surgery recovery process with your doctor. Potential risks include infection, pain, implant loosening, and loss of correction due to improper bone healing. For more information about risks, see the **Patient Risk Information**.

For information about how we handle your information, see our **Privacy Policy**.

LAPIPLASTY® and "ADDUCTOPLASTY®" are registered trademarks of Treace Medical Concepts, Inc

M2076B

