

TWICKENHAM PEDIATRICS

CHECK-UP & IMMUNIZATION SCHEDULE



This immunization schedule is recommended by the American Academy of Pediatrics.

Birth	In Hospital: 1st PKU and Hep B #1 (hospital may use one of several brands)
3-5 days	1 st Well Child Check
2 weeks	Well Child Check, 2nd PKU and ENGERIX-B (Hep B #1) if not given at birth
2-month	Well Child Check, Vaxelis (DTaP, IPV, HIB, and Hep B), Vaxneuvance (PCV-15), Rotateq
4-month	Well Child Check, Vaxelis (DTaP, IPV, HIB, and Hep B), Vaxneuvance (PCV-15), Rotateq
6-month	Well Child Check, Vaxelis (DTaP, IPV, HIB, and Hep B), Vaxneuvance (PCV-15), Rotateq
9-month	Well Child Check and Hemoglobin Level
12-month	Well Child Check, MMR, VARIVAX (Varicella), VAQTA (Hep A #1) and Lead Level
15-month	Well Child Check, DAPTACEL (DTaP), PedvaxHIB (HIB #4), Vaxneuvance (PCV-15)
18-month	Well Child Check, VAQTA (Hep A #2) and Autism and developmental screening
2 years	Well Child Check and Lead Level
3 years	Well Child Check
4 years	Well Child Check, ProQuad (MMRV), Quadracel (DTaP, IPV)
5 years	Well Child Check
11 years	Well Child Check, Adacel (Tdap), Gardasil 9 (HPV9 #1) - MenQuadfi #1 (if HPV9 declined)
11 ½ years	Gardasil 9 (HPV9 #2), MenQuadfi #1 (if not given at 11-year check-up) - No shots needed here if HPV9 is declined.
12-15 years	Well Child Check, Gardasil 9 (HPV9) series if not completed - if started before age 15 years 2-dose schedule, if started at 15 years or older 3-dose schedule is required
16 years	Well Child Check, MenQuadfi #2, Gardasil 9 (HPV9) series if not completed - if started before age 15 years 2-dose schedule, if started at 15 years or older 3-dose schedule is required, Bexsero series (2-dose schedule)
17-18 years	Well Child Check, MenQuadfi # 2 if not given at 16, Gardasil 9 (HPV9) series if not completed - if started before age 15 years 2-dose schedule, if started at 15 years or older 3-dose schedule is required, Bexsero series (2-dose schedule) if not previously given

* A Well Child Check is recommended every year after age 2.

** Annual Flu Vaccination is recommended for ages 6-months and up. Children up to age 8 receiving the flu vaccine for the first time must have 2 doses spaced at least 28 days apart. Flu vaccine brand may vary.