ZINC 9%

GUARANTEED ANALYSIS

Derived from: Zinc EDTA

WARNING: KEEP OUT OF REACH OF CHILDREN

May be harmful if swallowed. Causes mild skin irritation. May be harmful if inhaled. Causes eye irritation.

GENERAL INFORMATION

ZINC 9% is a EDTA chelated zinc solution designed for use in most soil applied liquid fertilizers and foliar sprays. ZINC 9% is intended for use and is compatible with phosphate fertilizers. This material should only be used as recommended. It may prove harmful when misused.

APPLICATIONS & RATES

SOIL APPLICATION: Use at the rate of 1 pint to 2 quarts per acre. Apply in a band at planting time with liquid fertilizer, knifed in preplant, or at sidedressing, as a supplement to broadcast applications.

TECHNICAL DATA
Net Weight:

Specific Gravity

DIRECTIONS FOR USE

pH at 68°F

Freezing Point

Weight Per Gallon lbs/gal @68°F

10.85

1.30

6-8 32°F

This product is intended for the correction of zinc deficiencies. This product should be used on the basis of a soil and/or tissue analysis. To

crop advisor, university crop extension publication, or a trained

agronomist. The ideal rate of application will vary among fields.

rate, and placement are important for best results.

determine fertilizer application rates suitable for your geographical area

or the maximum allowable application rate per acre, consult a certified

depending on soil pH. organic matter, and crop. Application of timing.

FOLIAR APPLICATION: Use at a rate of 1-2 quarts per acre in a minimum of 20 gallons of water for ground application. For aerial application apply in a minimum of 5 gallons per acre.

HIGH RESPONSE CROPS INCLUDE: corn, beans, cotton, sweet corn, sorghum, popcorn, onions, spinach, rice

OTHER CROPS: soybeans, sugar beets, wheat, alfalfa, barley, oats, tobacco, carrots, cabbage, table beets, lettuce, tomatoes, radishes, squash, rye, clover, bermuda, coastal grasses, sudangrass, peanuts

FERTILIZER MIXING INSTRUCTIONS *Always perform a compatibility (jar) test with all tank mix ingredients. Mix As Follows:

- 1. Partially fill tank with fertilizer and start agitation
- 2. Add ZINC 9% to circulating tank mix
- 3. Add remaining fertilizer
- 4. Add other tank mix ingredients (pesticides, nutrients, or additives) after successful compati-bility (jar) testing
- 5. Maintain good agitation throughout the tank mix

FIRST AID

Avoid contact with eyes and skin. Avoid breathing fumes or spray mist. May be harmful if swallowed. Wash thoroughly after handling. Remove and wash contaminated clothing before reuse.

If in eyes: Immediately flush eyes with water for at least 15 minutes. Seek immediate medical attention if irritation occurs.

If on skin or clothing: Flush skin with water. If irritation occurs, seek immediate medical attention.

If swallowed: Give large amounts of water and induce vomiting by touching back of throat with finger unless unconscious. Seek immediate medical attention.

PERSONAL PROTECTIVE EQUIPMENT

Wear gloves and protective clothing. Wear eye protection (glasses, goggles or face shield).

STORAGE & DISPOSAL

Store product in original container. DO NOT store below 40 °F or above 120 °F. DO NOT reuse container. DO NOT contaminate water, food or feed by storage or disposal. Keep packaging tightly closed. Triple rinse or equivalent and dispose of container in accordance with local, state and federal regulations. Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapf

ME I C	ONTENTS:	
X	2.5 gallons (27.13 lb)	





888-679-5103 www.egebio.com PO Box 538 Minneola. KS 67865

