



Well Check and Recommended Vaccine Schedule

Newborn: Hepatitis B if not given at birth

2 weeks: No vaccines, *Draw Newborn Screen Lab in office.*

1 month: No vaccines

2 months: Vaxelis (Dtap, Hepatitis B, Haemophilus influenzae type B, Polio), Vaxneuvance (Pneumococcal 15), Rotateq

4 months: Vaxelis (Dtap, Hepatitis B, Haemophilus influenzae type B, Polio), Vaxneuvance (Pneumococcal 15), Rotateq

6 months: Vaxelis (Dtap, Hepatitis B, Haemophilus influenzae type B, Polio), Vaxneuvance (Pneumococcal 15), Rotateq

9 months: No vaccines

12 months: MMRII (MMR), Varivax (Varicella), Vaqta (Hepatitis A); *Sent for CBC and Lead Labs*

15 months: Pentacel (Dtap, Haemophilus influenzae type B, Polio), Vaxneuvance (Pneumococcal 15)

18 months: Vaqta (Hepatitis A)

24 months: No vaccines

30 months: No vaccines

3 years: No vaccines

4 years: ProQuad (MMR, Varicella), Quadracel (Dtap, Polio)

5 years: No vaccines

6 years: No vaccines

7 years: No vaccines

8 years: No vaccines

9 years: No vaccines; *Sent for Lipid Panel Lab*

10 years: No vaccines

11 years: Menquadfi (Meningitis ACWY), Adacel (Tdap)

12 years: No vaccines

13 years: No vaccines

14 years: No vaccines

15 years: No vaccines

16 years: Menquadfi (Meningitis ACWY)

17 years: No vaccines

18 years: No vaccines; *Sent for Lipid Panel, HIV and HepC Labs*

Recommended vaccines (Not required by schools in Texas):

Gardasil (Human papilloma Virus): 2 dose series that can be started as early as 9 years old.

Bexsero (Meningitis B): 2 dose series that can be started as early as 16 years old.

Beyfortus (RSV): 1 dose series given either at birth or before the first RSV of life if mother did not receive vaccine 2 weeks prior to delivery.

Flu: Given annually. If younger than 9 years when receiving flu vaccine for first time, child will require a second dose within the same flu season 1 month apart. Subsequent doses are one annually.

COVID: See recommendations by manufacturer.